KEKD/ ^ /^ h^^/KE W W Z îîXìí/^^E ííóõrôôîï ~Kvo]v •
: vµ ŒÇ îìîî

3UR[LPLW\ WR KHDOWK FDUH FHQWUHV DQG V

&RPPXQLW\ &OLQLFV LQ %DQJODGHVK

6\HG +DVDQXO ,VODP ,QVWLWXWH RI +HDOWK 6FLHQFHV

6FKRRO RI (FRQRPLFV DQG)LQDQFH 0DVVH\ 8QLYHUVLW

dZ]• • Œ] • }vš]v• Á}Œl ^vZ‰ Œ}P Œ}v•}uš šZv &]v v U D }uÇvšv]À Œ•]šÇ
v Œ]š]]•u Œ]vÀ]š X Yµ}š š]}v• u Ç u }v Æ‰o]]š ‰ Œu]••]}v }

^Ç , • v
^ Z}}o }(}v}u] • v &]v v
D •• Ç hv]À Œ•]šÇ
WŒ]À š P ííîîî
W ou Œ•š}v E}ŒšZ ðð ðî
E t • > E
u]o^W X, • v›u •• ÇX XvÌ
WZ}v W =òð ò ïñò õìõõ Æšv ôðìíõ

Proximity to health care centres and service intake: The case of Community Clinics in Bangladesh*

Syed Hasan[y], Tasnima Akter[z], Musharrat Jahan[x], and Ashraf Dewan[{]

[y]School of Economics and Finance, Massey University, New Zealand

[z]Faculty of Health, University of Technology Sydney, Australia

[x]UNICEF, Bangladesh

[{] School of Earth and Planetary Sciences, Curtin University, Australia

January 19, 2022

Abstract

We investigate how distance from heath centres a ects service intake for women and children. Relying on ve rounds of recent nationally representative demographic and health survey data from Bangladesh, our logistic regression analyses reveal that proximity to health centres barely a ects the intake of health care services for women and children, even in the country's rural areas. Interestingly, this indicates that the newly established Community Clinics have not signi cantly contributed to the country's intake of health care services. The low service intake may result from

1. Introduction

The availability of health care service delivery centres (health centres hereafter) a ects the intake of their services (Abbott et al., 2017; Lu and Slusky, 2019; Lindo et al., 2020; De Luca et al., 2021; Abu-Qarn and Lichtman-Sadot, 2022). Previous studies nd take-up of health services to be sensitive to distances to the health centres (Kremer and Glennerster, 2011). For example, Lindo et al. (2020) nd substantial e ects of travel distances on abortion rates. Lu and Slusky (2019) nd driving distances to the nearest clinics to increase fertility rates. Thus, policies that reduce travel distances and travel times are likely to increase utilisation of health services (Karra et al., 2017).

The positive e ects of the proximity to health centres on service intake are likely to be higher in low-income settings where women and children with low socioeconomic backgrounds cannot access the services for infrastructural, religious, societal or nancial constraints (Ensor and Cooper, 2004; Jacobs et al., 2012; Raut and Tanaka, 2021; McGuire et al., 2021). For example, McGuire et al. (2021) nd distances to signi cantly reduce the probability of having a facility delivery in Malawi, which is particularly strong for women with lower socioeconomic status. Even relatively small distances from health facilities are associated with substantial mortality penalties for children in low- and middle-income countries (Karra et al., 2017). Thus, nding out the optimal location of health centres can improve access and ensure the best possible health outcome for a country (Adhvaryu and Nyshadham, 2015).[1]

While improving access to the formal health care sector is a primary public health goal in many low-income countries, the returns to the access are unclear as the quality of care at public health facilities is often considered inadequate (Adhvaryu and Nyshadham, 2015). Previous studies report improved outcomes from improving the quality of service delivery (Arifeen et al., 2005; Chowdhury et al., 2008; Hoque et al., 2014).[2] Thus, the quality of the o ered and/or delivered services can a ect the intake of health services. In particular, high-quality services may attract many stakeholders to

[1] It is possible to increase access to some types of health care services through home delivery (Herrera-Almanza and Rosales-Rueda, 2020). However, it is not appropriate for many kinds of services like ANC or referrals for fever.

[2] For example, Arifeen et al. (2005) nd improvement in the care quality of sick under- ve children in rst-level facilities in Bangladesh can improve their health outcomes. Chowdhury et al. (2008) nd appropriate training and supervision allow safe and e ective management of severe pneumonia. Hoque et al. (2014) nd regular supervision of workers, even with minimal pre-service training, improves the quality of child health care in rst-level health facilities.

enjoy them while people may take the services from other sources or even avoid them when their (or perceptions of the) qualities are poor.

Against this background, we primarily investigate how distances from the nearest health centres a ect service intake for women and children in Bangladesh. We also examine whether the service quality of some basic health centres, compared to the other types of health centres in the country, can be an issue. We choose to investigate the case of Bangladesh for two important reasons. First, the availability of several rounds of large scale demographic and health survey data includes the information of distances from di erent types of health care centres. Second, it is particularly interesting to investigate the contribution of the newly established/revitalised Community Clinics, on which the country is relying heavily for the future delivery of health care services (Planning Commiss(care)-4200aThl2.92 2020

Bangladesh has progressed impressively on di erent health indicators in the last two decades, compared to its' neighbours and many other developing countries (Planning Commiss(care)-4200aThl2.92 2020 Sachs et al.re 2021re The government aims to continue with the pace and achieve the targets of Sustainable Development Goals (SDGs) by the end of 2030 through strengthening the Community Based Health Care (CBHC) system. Along that line, Bangladesh has established more than one thousand Community Clinics (CCs), the agship program of the present government, to set up one centre for every 6,000 people (Planning Commiss(care)-4200aThl2.92 2020 The plan is that Community Clinics will assume the full responsibility of health population and nutrition of the entire population of its catchment area (Planning Commiss(care)-4200aThl2.92 2020

The government planned to establish the Community Clinics in 1996 to extend primary health care at the doorsteps of the villagers all over the country. The construction of Community Clinics started in 1998 and, during 1998-2001, about 10,723 of them were constructed, of which about 8,000 started functioning. The Community Clinics were operational only for a short time as they were closed in 2001 after the change of the government and remained closed till 2008. The government planned to revitalize the Community Clinics in 2009. Since then, around 13,831 facilities have been constructed, with 4,000 more clinics to be built by the end of 2022 (Planning Commiss(care)-4200aThl2.92 2020

]TJ 091 Tf 455.709 0 Td f 45fE (

illness, including nutritional education and micronutrient supplements, iii) screening and referrals for noncommunicable diseases such as hypertension, diabetes, arsenicosis, cancer, heart disease and autism, iv) treatment of minor ailments and rst aid for simple injuries, v) health and family planning education and counselling and identi cation of emergency and complicated cases with referrals and vi) free of cost essential medicines. The Community Clinics provide a woman-friendly platform where about 95 per cent of service seekers are women and children availing various primary health care services near their home (Planning Commission, 2020

2. Method

We have employed the following speci cation to model an individual's likelihood to intake a particular health care service:

$$y_i = \quad + \quad D_i + \quad X_i + \quad _s + \quad _t + \quad "_i ; \qquad (1)$$

where, for each individual i, the outcome variable y takes the value of one if someone has taken a particular health care service and zero otherwise. We use separate regressions to model the intake of three crucial health care services for reproductive-age women { i) use of modern birth control instruments, ii) making four or more ANC visits and iii) checking up by a trained provider within 42 days of delivery. We also use separate regressions for the intake of three services by the children{i) having all vaccines (for children aged 12-23 months), ii) receiving vitamin A in the last six months (for children aged 9-59 months) and iii) taking advice or treatment for fever symptoms (for under- ve children). The reason for choosing these dependent variables is that they are important health indicators and Community Clinics (and some urban health care centres) provide services in those health areas to women and children.[3]

The exposure variableS9091 Tfw9 -J/Ft sn -Tdvicetake7(mon)01r7(enden)28(t)-3

a series of categorical/dummy variables a ecting the health service intake of the stakeholders. The analysis with women includes a person's age, own education, partner's education, socio-economic status, and rural/urban location of residence. For our investigation with children, the vector X additionally includes the sex of the child and child's age. Note that women's age and education and partner's education, used in the analysis for women, are replaced by the same categories for mother and father.

The term , division xed e ects, accounts for possible omitted location variables and the time-invariant di erences in administrative divisions a ecting service intake. We also include , survey rounds xed e ects, to capture the changes in the service delivery over the years. Finally," is the independently and identically distributed error term.

The problem with the above models is that the variable `distance from the nearest health centre' may su er from endogeneity. The potential source of endogeneity in our model may engender from three sources. First, the distribution of health care centres is non-random. It can be the case that the government has established the health centres at locations with higher demand for its services. Second, residential sorting, in which high service intake households may choose to relocate near the health centres to enjoy the bene t of health care services permanently. Third, greater employment opportunities near the health care centres may attract young people who may have a higher demand for some health care services. For example, they can take birth control measures in a higher proportion as they may have a lower desire to have children. All of the factors are likely to be negatively associated with distance from health centres, thus making the exposure variable endogenous to the models for health care services. Not controlling for those factors in the empirical models will overestimate the average e ect of distance on the intake of health services.

Endogeneity in empirical models is typically addressed by employing the instrumental variable (IV) approach. Unfortunately, an important limitation of IV estimation is that it estimates the local average treatment e ect (LATE) rather than the average treatment e ect (ATE), a more policy-relevant quantity. In other words, the estimated impacts of treatments through the IV approach do not usually represent the case for the entire population, nor even to all of the treated observations. Instead, the IV estimate provides the treatment e ect only for the people whose choice of treatment

was a ected by the instrument. Thus, the estimated treatment e ects may vary with the choice of the instruments.[4]

On the other hand, the discussed sources of endogeneity in our models are less likely to be an issue in the context of Bangladesh. The low concern for endogeneity is because, rst, the health centres in the country are likely to be established with the consideration of equitable distributions across administrative regions, rather than the demand for health services (Planning Commission, 2020a). Second, individuals are less likely to relocate near the health centres since the marginal cost of relocation can be higher than the transportation (and time) cost of visiting the health centres occasionally. Finally, the employment opportunities created by the health care centres, especially when they are small, are likely to be trivial to attract the young people with higher demand for health care services. Therefore, we have not considered employing the IV approaches. However, recognising the endogeneity concern, we interpret the coe cients of the exposure variable as the upper bounds of the average treatment e ect of proximity to health centres.

We analyse ve rounds of Bangladesh Demographic and Health Survey (BDHS) data { 2004, 2007, 2011, 2014, and 2017. The BDHS is a cross-sectional survey collecting nationally represen-

ing weights to get a total of 73,040 women. Thus, our analysis of women's birth control service intake includes 11,440, 10,996, 17,749, 17,863, and 20,127 women for 2004, 2007, 2011, 2014, and 2017 rounds, respectively.

We follow similar restrictions for constructing samples for analysing other health service delivery for women. For the case of (four or more) ANC visits, restricting our sample to mothers with less than ve years of old children leaves us a total of 27,101 observations. From that, we drop 11 observations with missing weights to get the nal analysis sample of 27,090 women. For the analysis of the intake of PNC service by a trained provider, we additionally drop 1,161 mothers who either have not mentioned whether they have their health checked after hospital discharge (or home delivery) or fail to mention the type of the service provider. Thus our nal sample for the analysis of PNC service by a trained provider is 25,929 women.

Birth records data in the survey include a total of 201,576 children, of which 33,605, 30,527, 45,844, 43,772 and 47,828 belong to 2004, 2007, 2011, 2014 and 2017 survey rounds, respectively. From that, we only retain 7,223 last-born children aged 12-23 months to construct our analysis sample for child vaccination. On the other hand, for the analysis of vitamin A intake in the last six months, we retain 26,406 last-born children who belong to the age of 9-59 months. We then drop 963 observations for missing information about vitamin A intake and 2 observations for missing wights. Thus our analysis sample for vitamin A intake includes 25,441 children. Finally, for the analysis of advice sought for fever, we retain 11,755 under- ve children who have su ered from fever in the last two weeks of the survey. From that, we drop 4 observations for missing wights to get our analysis sample of 11,751 children[5].

The primary outcomes of interest are three di erent health service intakes each for women and children. Here, the three binary outcomes for women are de ned as follows: i) uses birth control methods, that takes a value of one if anyone currently uses a modern method and zero otherwise, ii) has made adequate ANC visits, that takes a value of one if anyone makes four or more ANC

The three important health services for children are de ned as follows: i) vaccination, in which children aged 12-23 months who have received all vaccines takes a value of one and zero otherwise, ii) vitamin A intake, in which child (aged 9-59 months) who has received vitamin A in the last six months takes a value of one and zero otherwise, and iii) have sought referral for fever, in which among under- ve children with fever symptoms who have received advice or treatment takes a value of one and zero otherwise. These health care services are important indicators of the usefulness of health facilities to the catchment population, and therefore we have employed them in our analysis.

Proximity to the nearest health facility is the main variable of interest in the study. We categorise the variable as dichotomous, taking a value of one when someone lives further than two kilometres of any health centre and zero otherwise. We choose the dummy variable due to its' straightforward policy implications. However, we have also discussed the consequence of using di erent forms of distances in the model to con rm that our results are not dependent on the choice of distance cuto . Table 1 presents year-wise summary statistics of the dependent variables, indicating signi cant di erences in some indicators between people living within two kilometres of health care centres against those living apart.

[Table 1]

Based on the previous literature, we consider a total of ve potential confounding factors in the analyses. For the analysis of women's health care services, they are woman's age (seven age groups: 15-19, 20-24, 30-34, 35-39, 40-44 and 45-49 years, with the 25-30 years group as the reference category), woman's education (primary, secondary and higher, with the uneducated women as the reference category), partner's education (primary, secondary and higher, with the uneducated partner as the reference category), socio-economic status (poorer, middle, richer and richest, with the poorest group as the reference category), rural (with the urban households as the reference category), administrative divisions (Barishal, Chattogram, Khulna, Rajshahi, Rangpur and Sylhet, with Dhaka as the reference category) and time (2007, 2011, 2014 and 2017 waves, with the 2004 wave as the reference category). For the analysis with children, we additionally include female (with the male as the reference category) and child's age (less than 12 months and 12-23 months, with the child aged 24 or more months as the reference category). Table 2 presents the number of

observations in each category of the exposure and control variables employed to analyse the health service intake of women and children in Bangladesh.

[Table 2]

3. Results and Discussion

We estimate model (1) for each of the dependent variables considered in this study. With the binary nature of our dependent variables, we estimate both the linear probability models (LPMs) and the logistic regression models but prefer the latter[6]. Each of the independent variables takes the form of a dummy variable, and so the OLS estimates and the marginal e ects (MEs) from the logit models indicate the percentage point (pp) changes in the probability of a service intake for a particular group/variable, compared to the reference group. Note that the reference groups for the analysis of woman health service intake are women living within two kilometres of any health centres, belong to the lowest SES group, has no education both for herself and her partner, located in the urban areas and the Dhaka division, aged 25-29 years and surveyed in the 2004 round of BDHS.

The LPM estimates in Column 1 of Table 3 indicates no signi cant e ect of distance on the use of modern contraceptives[7]. This is not surprising as many birth control devices are widely available in the local shops in Bangladesh, as we observe in the latest BDHS survey (NIPORT and ICFI, 2020). For all other control variables, we mostly observe signi cant e ects that are also in line with our expectations. For example, women's age demonstrates a quadratic relationship with the use of contraceptives, which remain lower at a young age and then increase until they reach 35-39 years and decrease after that. The pattern is generally expected as women's demand for contraceptives increase until they reach a certain age, which then falls as they move towards the end of their reproductive age. Previous studies nd that many women, believing that fertility declines from mid-30s, stop using contraception once they reach 40 years of age (Allen et al., 2013).

[Table 3]

6

Own schooling generally increases contraceptive use. Better-educated women are likely to have preferences for smaller families and thus higher demand for contraceptives (Di Novi and Thakare, 2020). On the other hand, women with partners having secondary or higher education have a lower use for contraceptives. The observed pattern of birth control measures can be due to an increased use of birth control devices for men, associated with their increase in education, as we observe in BDHS (NIPORT and ICFI, 2020). Interestingly, fewer women in the topmost socioeconomic status (SES) group use contraceptives compared to the bottom one. Some previous studies nd a similar pattern and explain it as a consequence of the national family planning programs that usually target poor women (Vu et al., 2016).

There are regional variations in the use of contraceptives, although rural women use them to a lesser extent, as observed in earlier research (Di Novi and Thakare, 2020). Fewer women in two conservative regions in Bangladesh {Chittagong (Chottogram) and Sylhet divisions {use modern birth control measures compared to their counterparts in the Dhaka division; more women in the remaining four divisions use modern contraceptives. Finally, the use of modern birth control methods increase over the survey periods. The overtime increase in modern contraceptive use is likely due to the government encouragement for family planning as well as the overall modernisation of the country (UNDP, 2017). All these results are consistent with some previous studies. For example, Amin et al. (2010) nd that mother's education, wealth index and place of residence are closely associated with access to maternal and newborn healthcare services in Bangladesh.

The marginal e ects from the logit model in Column 2 are comparable to the OLS estimates. Since this generally holds for our entire analysis, we will only consider the former to explain the intake of other health care services.

Column 4 presents the results from the model of taking (at least four) ANC visits. The results indicate that the ANC service intake is lower by 4.1 percentage points for the people living further than two kilometres of any health centres, compared to their counterparts living within two kilometres of the centres. The negative e ect of distance is a re ection of the fact that pregnant women

higher birth orders that may reduce the need for antenatal care services. Akter et al. (2016) nd women with more than one child utilising ANC services to a lower extent, due to their previous pregnancy experiences and time constraint resulting from an increased responsibility of child care. On the other hand, higher education and SES can increase the demand for the service among women (Di Novi and Thakare, 2020). Again rural mothers take lower ANC services, but the intake varies by administrative divisions, re ecting the di erence in socio-cultural factors (Di Novi and Thakare, 2020). Finally, the ANC service intake increases over time which is due to the increased awareness for maternal and child health care services in Bangladesh (Akter et al., 2018).

Next, we discuss the model of PNC service intake from a trained provider within 42 days of delivery. Column 6 results indicate that the postnatal care service intake for the mothers living further than two kilometres of any health centres is lower than their counterparts living within two kilometres of the centres. In particular, the PNC service intake is lower by 3.5 percentage points for the former than the latter group, but the e ect is not statistically signi cant. The results for all other control variables closely follow that of the ANC intake.

For the analysis of the intake of health services by children, the reference groups are boys older than two years, have uneducated parents, belong to the lowest SES group, lives within two kilometres from any health centres, located in the urban areas and Dhaka division, whose mother's age is 25-29 years and are surveyed during the 2004 round of BDHS. Our results in Column 8 indicate that the probability of having (all the required) vaccination is 3.8 percentage points lower for children living over two kilometres of any health centres compared to their counterparts, but the e ect is not statistically signi cant. MEs of other control variables indicate a lower vaccine intake for a child with older mothers (who are usually unaware), but as observed in (Di Novi and Thakare, 2020), the intake increases with parental education and wealth, indicating the role of consciousness and capacity in vaccination. Finally, we observe signi cant variation in the vaccination completion rates among children of di erent regions and periods.

Results from the model of Vitamin A intake, presented in Column 10, indicate a statistically signi cant and negative impact of living further from the health centres. In particular, the results indicate that the Vitamin A intake is signi cantly lower by 5.2 percentage points for children living two kilometres further from health centres, compared to the reference group. The e ect of other control variables in the model, for similar reasons, closely follow the results for vaccination.